



Anti-DPP4 monoclonal antibody, clone 22E8 (DCABH-432)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	Mouse monoclonal to CD26
Antigen Description	Cell surface glycoprotein receptor involved in the costimulatory signal essential for T-cell receptor (TCR)-mediated T-cell activation. Acts as a positive regulator of T-cell coactivation, by binding at least ADA, CAV1, IGF2R, and PTPRC. Its binding to CAV1 and CARD11 induces T-cell proliferation and NF-kappa-B activation in a T-cell receptor/CD3-dependent manner. Its interaction with ADA also regulates lymphocyte-epithelial cell adhesion. In association with FAP is involved in the pericellular proteolysis of the extracellular matrix (ECM), the migration and invasion of endothelial cells into the ECM. May be involved in the promotion of lymphatic endothelial cells adhesion, migration and tube formation. When overexpressed, enhanced cell proliferation, a process inhibited by GPC3. Acts also as a serine exopeptidase with a dipeptidyl peptidase activity that regulates various physiological processes by cleaving peptides in the circulation, including many chemokines, mitogenic growth factors, neuropeptides and peptide hormones. Removes N-terminal dipeptides sequentially from polypeptides having unsubstituted N-termini provided that the penultimate residue is proline.
Immunogen	Recombinant full length Human CD26 produced in HEK293T cells (NP_001926)
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	Human
Clone	22E8
Purification	Purified from mouse ascites fluids by affinity chromatography
Conjugate	Unconjugated

Applications	WB, IHC-P, Flow Cyt, ICC/IF
Positive Control	IHC-P: Liver tissue Flow Cyt: Hela cells WB: Transfected HEK293T cell lysate ICC/IF: Transfected COS7 cells
Format	Liquid
Size	100 µl
Buffer	pH: 7.30; Preservative: 0.02% Sodium azide; Constituents: 48% PBS, 50% Glycerol, 1% BSA
Preservative	0.02% Sodium Azide
Storage	store at -20°C. Avoid repeated freeze / thaw cycles.
Ship	Shipped at 4°C.

GENE INFORMATION

Gene Name	DPP4 dipeptidyl-peptidase 4 [Homo sapiens]
Official Symbol	DPP4
Synonyms	DPP4; dipeptidyl-peptidase 4; ADCP2, adenosine deaminase complexing protein 2 , CD26, dipeptidylpeptidase IV (CD26, adenosine deaminase complexing protein 2); dipeptidyl peptidase 4; DPPIV; ADCP-2; DPP IV; dipeptidylpeptidase 4; dipeptidyl peptidase IV;
Entrez Gene ID	1803
Protein Refseq	NP_001926
UniProt ID	P27487
Chromosome Location	2q23-qter
Pathway	Incretin Synthesis, Secretion, and Inactivation, organism-specific biosystem; Integration of energy metabolism, organism-specific biosystem; Metabolism, organism-specific biosystem; Protein digestion and absorption, organism-specific biosystem; Protein digestion and absorption, conserved biosystem; Regulation of Insulin Secretion, organism-specific biosystem; Synthesis, Secretion, and Inactivation of Glucagon-like Peptide-1 (GLP-1), organism-specific biosystem;
Function	aminopeptidase activity; collagen binding; dipeptidyl-peptidase activity; peptidase activity; peptide binding; protease binding; protein binding; protein homodimerization activity; receptor activity; receptor binding; serine-type endopeptidase activity; s