

Anti-HSV type 2 Monoclonal antibody, Clone A911 (DMAB3603)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Specificity	HSV2, specific for gD.
Target	HSV type 2
Immunogen	Purified HSV-2 virions
Isotype	IgG1
Source/Host	Mouse
Species Reactivity	HSV
Clone	A911
Affinity Constant	Not determined
Purification	90% pure. Protein A chromatography
Conjugate	Unconjugated
Applications	Suitable for use in Western blot, ELISA and IFA. Each laboratory should determine an optimum workingtiter for use in its particular application. Other applications have not been tested but use in such assays shouldnot necessarily be excluded.
Format	Purified, Liquid
Concentration	100ug/ml (OD280nm, E0.1% = 1.3)
Size	100 μg
Buffer	0.01M PBS, pH 7.2. Product contains no stabilizing proteins.

45-1 Ramsey Road, Shirley, NY 11967, USA

© Creative Diagnostics All Rights Reserved

Tel: 1-631-624-4882 Fax: 1-631-938-8221

Storage

Upon receipt, aliquot and store at -20°C. Avoid multiple freeze/thaw cycles.

BACKGROUND

Introduction	Herpes simplex type 2 (HSV2) belongs to a familythat includes HSV1, Epstein-Barr virus (EBV) and Varicella zoster (chickenpox) virus. HSV1 and HSV2 are extremely difficult to distinguish from eachother. These viruses have a DNA genome, an icosahedral protein coat and areencased in a lipid membrane derived from the nuclear membrane of the lasthost. These viruses are capable of entering a latent phase where the hostshows no visible sign of infection and levels of infectious agent become verylow. During the latent phase the viral DNA is integrated into the genome of the host cell.
Keywords	Herpesviridae; Alphaherpesvirinae; Simplexvirus; Herpes simplex virus 2; HSV 2; Herpes Simplex Virus Type 2; HSV-2; Glycoprotein D precursor; Herpes Simplex Virus Type 2 Glycoprotein D; HHV2gp69; Herpesvirus 2; US6; Virion glycoprotein D; Herpes Simplex Virus Type 2 Glycoprotein D; HSV-2 gD

© Creative Diagnostics All Rights Reserved