

## Anti-CDK5 (aa 229-292) polyclonal antibody (DPABH-03356)

This product is for research use only and is not intended for diagnostic use.

## **PRODUCT INFORMATION**

Antigen Description	Probably involved in the control of the cell cycle. Interacts with D1 and D3-type G1 cyclins. Can phosphorylate histone H1, tau, MAP2 and NF-H and NF-M. Also interacts with p35 which activates the kinase.
Immunogen	Synthetic peptide, corresponding to a region within C terminal amino acids 229-292 of Human Cdk5.
Isotype	IgG
Source/Host	Rabbit
Species Reactivity	Mouse, Human
Purification	Immunogen affinity purified
Conjugate	Unconjugated
Applications	WB, IHC-P, ICC/IF
Format	Liquid
Size	100 μΙ
Buffer	pH: 7.00; Constituents: 0.75% Glycine, 1.21% Tris, 10% Glycerol
Preservative	None
Storage	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## **GENE INFORMATION**

Tel: 1-631-624-4882 Fax: 1-631-938-8221

<sup>45-1</sup> Ramsey Road, Shirley, NY 11967, USA

Gene Name	CDK5 cyclin-dependent kinase 5 [ Homo sapiens ]
Official Symbol	Cdk5
Synonyms	CDK5; cyclin-dependent kinase 5; PSSALRE; TPKII catalytic subunit; protein kinase CDK5 splicing; cell division protein kinase 5; serine/threonine-protein kinase PSSALRE; tau protein kinase II catalytic subunit;
Entrez Gene ID	<u>1020</u>
Protein Refseq	<u>NP_001157882</u>
UniProt ID	<u>Q00535</u>
Chromosome Location	7q36
Pathway	Alzheimers disease; Axon guidance; CRMPs in Sema3A signaling; Cocaine addiction;
Function	ATP binding; ErbB-2 class receptor binding; ErbB-3 class receptor binding; acetylcholine receptor activator activity; cyclin-dependent protein kinase activity; kinase activity; nucleotide binding