



## Anti-Goat IgG polyclonal antibody [Biotin] (DPBT-67075DG)

This product is for research use only and is not intended for diagnostic use.

### PRODUCT INFORMATION

<b>Product Overview</b>	Donkey Anti Goat IgG (H/L),BiotinDonkey Anti Goat IgG (H/L),Biotin
<b>Immunogen</b>	Goat IgG
<b>Isotype</b>	IgG
<b>Source/Host</b>	Donkey
<b>Species Reactivity</b>	Goat
<b>Conjugate</b>	Biotin
<b>Applications</b>	IHC, ELISA
<b>Format</b>	Purified IgG conjugated to Biotin - liquid
<b>Concentration</b>	0.5 mg/ml
<b>Size</b>	1 mg
<b>Buffer</b>	Phosphate buffered saline
<b>Preservative</b>	0.1% Sodium Azide
<b>Storage</b>	Store at +4 °C or at -20 °C if preferred.Storage in frost-free freezers is not recommended.This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

### BACKGROUND

**Introduction**

Immunoglobulin G (IgG) are antibody molecules. Each IgG is composed of four peptide chains - two heavy chains  $\gamma$  and two light chains. Each IgG has two antigen binding sites. Other Immunoglobulins may be described in terms of polymers with the IgG structure considered the monomer. IgG molecules are synthesized and secreted by plasma B cells. IgG antibodies are large molecules of about 150 kDa composed of 4 peptide chains. It contains 2 identical heavy chains of about 60kDa and 2 identical light chains of about 25 kDa, thus a tetrameric quaternary structure. The two heavy chains are linked to each other and to a light chain each by disulfide bonds. The resulting tetramer has two identical halves, which together form the Y-like shape. Each end of the fork contains an identical antigen binding site. The Fc regions of IgGs bear a highly conserved N-glycosylation site. The N-glycans attached to this site are predominantly core-fucosylated diantennary structures of the complex type. In addition, small amounts of these N-glycans also bear bisecting GlcNAc and  $\alpha$ -2,6-linked sialic acid residues.

**Keywords**

Ig gamma 1 chain C region; IGHG1; Immunoglobulin heavy constant gamma 1; Immunoglobulin G; IgG; IgG heavy chain; Immunoglobulin G heavy chain